

3067

NDC 59556-764-13

# Ibuprofen Capsules

## 200 mg

Solubilized Ibuprofen Capsules, 200 mg  
Pain Reliever/Fever Reducer (NSAID)

80 Liquid Filled Capsules

Strides Pharma Inc.

This package for households without young children

Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.

**Drug Facts**

Active ingredient (in each capsule)	Purpose
Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)* (present as the free acid and potassium salt)	Pain reliever/ Fever reducer

\*nonsteroidal anti-inflammatory drug

**Uses**

- temporarily relieves minor aches and pains due to:
  - headache
  - backache
  - the common cold
  - toothache
  - menstrual cramps
  - muscular aches

Manufactured by:  
**Strides Shasun Limited**  
Bengaluru - 560076, India

Distributed by:  
**Strides Pharma Inc.**  
East Brunswick, NJ 08816

Revised: 08/2017

Code No.:  
KR/DRUGS/KTK/25/415/98

LOT: \_\_\_\_\_

Exp: \_\_\_\_\_

NO VARNISH ZONE

PEEL HERE

1031853

Dimension : 150 x 50 mm

<p><b>Drug Facts (continued)</b></p> <ul style="list-style-type: none"> <li>■ minor pain of arthritis</li> <li>■ temporarily reduces fever</li> </ul> <p><b>Warnings</b></p> <p><b>Allergy alert:</b> Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:</p> <ul style="list-style-type: none"> <li>■ hives</li> <li>■ asthma (wheezing)</li> <li>■ skin reddening</li> <li>■ facial swelling</li> <li>■ shock</li> <li>■ rash</li> <li>■ blisters</li> </ul> <p>If an allergic reaction occurs, stop use and seek medical help right away.</p> <p><b>Stomach bleeding warning:</b> This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you</p> <ul style="list-style-type: none"> <li>■ are age 60 or older</li> <li>■ have had stomach ulcers or bleeding problems</li> <li>■ take a blood thinning (anticoagulant) or steroid drug</li> </ul>	<p><b>Drug Facts (continued)</b></p> <ul style="list-style-type: none"> <li>■ take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]</li> <li>■ have 3 or more alcoholic drinks every day while using this product</li> <li>■ take more or for a longer time than directed</li> </ul> <p><b>Heart attack and stroke warning:</b> NSAIDs, except aspirin, increase the risk of heart attack, heart failure and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.</p> <p><b>Do not use</b></p> <ul style="list-style-type: none"> <li>■ if you have ever had an allergic reaction to any other pain reliever/fever reducer</li> <li>■ right before or after heart surgery</li> </ul> <p><b>Ask a doctor before use if</b></p> <ul style="list-style-type: none"> <li>■ stomach bleeding warning applies to you</li> </ul>	<p><b>Drug Facts (continued)</b></p> <ul style="list-style-type: none"> <li>■ you have problems or serious side effects from taking pain relievers or fever reducers</li> <li>■ you have a history of stomach problems, such as heartburn</li> <li>■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma or had a stroke</li> <li>■ you are taking a diuretic</li> </ul> <p><b>Ask a doctor or pharmacist before use if you are</b></p> <ul style="list-style-type: none"> <li>■ under a doctor's care for any serious condition</li> <li>■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin</li> <li>■ taking any other drug</li> </ul> <p><b>When using this product</b></p> <ul style="list-style-type: none"> <li>■ take with food or milk if stomach upset occurs</li> </ul> <p><b>Stop use and ask a doctor if</b></p> <ul style="list-style-type: none"> <li>■ you experience any of the following signs of stomach bleeding:</li> </ul>
--	---	---

<p><b>Drug Facts (continued)</b></p> <ul style="list-style-type: none"> <li>■ feel faint</li> <li>■ have bloody or black stools</li> <li>■ have stomach pain that does not get better</li> <li>■ you have symptoms of heart problems or stroke:             <ul style="list-style-type: none"> <li>■ chest pain</li> <li>■ weakness in one part or side of body</li> <li>■ slurred speech</li> <li>■ leg swelling</li> </ul> </li> <li>■ pain gets worse or lasts more than 10 days</li> <li>■ fever gets worse or lasts more than 3 days</li> <li>■ redness or swelling is present in the painful area</li> <li>■ any new symptoms appear</li> </ul> <p><b>If pregnant or breast-feeding,</b> ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery</p>	<p><b>Drug Facts (continued)</b></p> <p><b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away</p> <p><b>Directions</b></p> <ul style="list-style-type: none"> <li>■ do not take more than directed</li> <li>■ the smallest effective dose should be used</li> <li>■ adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist</li> <li>■ if pain or fever does not respond to 1 capsule, 2 capsules may be used</li> <li>■ do not exceed 6 capsules in 24 hours, unless directed by a doctor</li> <li>■ children under 12 years: ask a doctor</li> </ul> <p><b>Other information</b></p> <ul style="list-style-type: none"> <li>■ each capsule contains: potassium 17.840 mg</li> <li>■ read all warnings and directions before use. Keep carton.</li> </ul>	<p><b>Drug Facts (continued)</b></p> <ul style="list-style-type: none"> <li>■ store at 20-25°C (68-77°F)</li> <li>■ avoid excessive heat above 40°C (104°F)</li> </ul> <p><b>Inactive ingredients</b> FD&amp;C blue 1, gelatin (hydrolyzed), medium chain triglyceride, PEG 600, potassium hydroxide, sorbitan and sorbitan solution. Printing ink: white edible oil, propylene glycol, shellac resins, sodium lauryl sulphate, isopropyl alcohol, titanium dioxide.</p> <p><b>Questions or comments?</b> call toll free 1-877-244-9825</p> <p style="text-align: center;">For most recent product information Visit us at <a href="http://www.stridesshasun.com">www.stridesshasun.com</a></p> <p>Manufactured by: <b>Strides Shasun Limited</b> Bengaluru - 560076, India.</p> <p>Distributed by: <b>Strides Pharma Inc.</b> East Brunswick, NJ 08816</p>
---	--	---

GLUE AREA  
Do not print

COLOUR:  Pantone 3272 C  Pantone 340 C  Pantone 317 C  Pantone 877 C  Black

Labeling element	font Size
Drug Facts heading	7 points
Drugs Facts	7 points
Heading	7 points
Sub heading	6 points
Bulleted Text	6 points
Leading	0.5 points
Bullets	6 points

Dimension : 150 x 50 mm

NOTE: Label dimension to be change during commercial time